



# ***Equioxx***<sup>®</sup>

## ***Injection*** (firocoxib)

**Non-steroidal anti-inflammatory drug  
for intravenous use in horses only.**

### **Information for Horse Owners**

**Indication:** EQUIOXX Injection is administered for up to 5 days for the control of pain and inflammation associated with osteoarthritis in horses. If further treatment is needed, EQUIOXX Oral Paste can be used for up to an additional 9 days of treatment. The overall duration of treatment with EQUIOXX Injection and EQUIOXX Oral Paste will be dependent on the response observed, but should not exceed 14 days.

This summary contains important information about EQUIOXX. You should read this information before you start giving your horse EQUIOXX and review it each time your prescription is refilled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or you want to know more about EQUIOXX.

---

#### **What is EQUIOXX?**

EQUIOXX is a veterinary prescription non-steroidal anti-inflammatory drug (NSAID) of the coxib class used to control pain and inflammation associated with osteoarthritis in horses. Osteoarthritis (OA) is a painful condition caused by progressive "wear and tear" of cartilage and other parts of the joints that may result in the following changes or signs in your horse:

- Limping or lameness.
- Decreased activity or exercise (reluctance to stand, walk, trot or run, or difficulty in performing these activities).
- Stiffness or decreased movement of joints.

---

#### **How to give EQUIOXX to your horse?**

EQUIOXX should be given according to your veterinarian's instructions. Do not change the way you give EQUIOXX to your horse without first

speaking with your veterinarian. Do not exceed 14 days of treatment. The recommended dosage of EQUIOXX Injection for intravenous administration in horses is 0.04 mg/lb (0.09 mg/kg) of body weight once daily for up to 5 days. If further treatment is needed, EQUIOXX (firocoxib) Oral Paste for horses can be used at a dosage of 0.045 mg/lb (0.1 mg/kg) bodyweight for up to an additional 9 days of treatment. The overall duration of treatment with EQUIOXX Injection and EQUIOXX Oral Paste will be dependent on the response observed, but should not exceed 14 days.

---

### **What kind of results can I expect when my horse is on EQUIOXX for OA?**

While EQUIOXX is not a cure for osteoarthritis, it can control the pain and inflammation associated with OA and can improve your horse's mobility.

- Response varies from horse to horse, but improvement can be quite dramatic.
  - Improvement can be seen in just a few hours in most horses.
- 

### **Which horses should not receive EQUIOXX?**

Your horse should not be given EQUIOXX if he/she:

- Has an allergic reaction to firocoxib, the active ingredient in EQUIOXX.
  - Has previously had an allergic reaction (such as hives, facial or lower limb swelling, or red or itchy skin) to aspirin or other NSAIDs.
  - Is presently taking aspirin, phenylbutazone, flunixin meglumine, diclofenac, ketoprofen, or other NSAIDs or corticosteroids.
  - The safety of EQUIOXX has not been determined in horses less than one year of age or in breeding horses, pregnant or lactating mares.
- 

### **EQUIOXX Injection should only be given intravenously to horses.**

- EQUIOXX is not for use in horses intended for human food consumption.
  - People should not take EQUIOXX. Keep EQUIOXX and all medications out of the reach of children. Consult a physician in case of accidental ingestion by humans.
- 

### **What to tell/ask your veterinarian before giving EQUIOXX.**

Talk to your veterinarian about:

- The signs of OA you have observed in your horse, such as limping or stiffness.
- If any tests, such as X-rays, will be done before EQUIOXX is prescribed.

- How often your horse may need to be examined by your veterinarian.
- The risks and benefits of using EQUIOXX.

Tell your veterinarian if your horse has ever had the following medical problems:

- Any side effects from taking EQUIOXX or other NSAIDs, such as aspirin or phenylbutazone.
- Any kidney disease.
- Any liver disease.
- Any gastrointestinal ulcers.

Tell your veterinarian about:

- Other medical problems or allergies that your horse has now, or has had in the past.
- All medicines that you are giving or plan to give to your horse, including those you can get without a prescription and any dietary supplements.

Tell your veterinarian if you plan to breed your horse, or if your mare is pregnant or nursing a foal.

### **What are the possible side effects that may occur in my horse during EQUIOXX therapy?**

EQUIOXX, like other NSAIDs, may cause some side effects. Serious side effects associated with NSAID therapy in horses can occur with or without warning. The most common side effects associated with EQUIOXX therapy involve the tongue, lips and skin of the mouth and face (erosions and ulcers of the mucosa and skin) and the kidney. Gastrointestinal, kidney and liver problems have also been reported with other NSAIDs. Look for the following side effects that may indicate your horse is having a problem with EQUIOXX or may have another medical problem:

- Sores or ulcers on the tongue and inside of mouth.
- Sores, scabs, redness, or rubbing of the facial skin, particularly around the mouth.
- Change in eating or drinking habits (frequency or amount consumed).
- Change in urination habits (frequency or color).
- Yellowing of gums, skin, or whites of the eyes (jaundice).
- Unexpected weight loss.
- Change in behavior (such as increased or decreased activity level).

It is important to stop therapy and contact your veterinarian if you think your horse has a medical problem or side effect while taking EQUIOXX. If you have additional questions about possible side effects, talk with your veterinarian or call 1-877-217-3543.

### **Can EQUIOXX be given with other medications?**

EQUIOXX should not be given with other NSAIDs



(for example, aspirin, phenylbutazone, diclofenac, ketoprofen or flunixin) or systemic corticosteroids (for example, prednisone, cortisone, dexamethasone, or triamcinolone).

Tell your veterinarian about all medications that you have given your horse in the past, and any medications you are planning to give with EQUIOXX. This should include other medicines that you can get without a prescription or any dietary supplements. Your veterinarian may want to check that all of your horse's medicines can be given together.

---

### **What do I do in case my horse receives more than the prescribed amount of EQUIOXX?**

- Consult your veterinarian if your horse receives more than the prescribed amount of EQUIOXX.

---

### **What else should I know about EQUIOXX?**

- This sheet provides a summary of information about EQUIOXX paste and general information about NSAIDs. If you have any questions or concerns about EQUIOXX or osteoarthritis pain, talk with your veterinarian.
- As with all prescribed medicines, EQUIOXX should only be given to the horse for which it is prescribed. It should be given to your horse only for the condition for which it is prescribed, at the labeled dose and duration.
- It is important to periodically discuss your horse's response to EQUIOXX. Your veterinarian will determine if your horse is responding as expected and if your horse should continue receiving EQUIOXX.

Manufactured for:

Merial Limited

Duluth, GA 30096-4640, U.S.A.

1-877-217-3543

Made in Germany

NADA 141-313, Approved by FDA

®EQUIOXX is a registered trademark of Merial.

©2011 Merial. All Rights Reserved.

Rev. 10-2011

722175 US2

